

**How I Found My Place
in the Universe
(and helped everybody else find theirs)**

**By
Jessica Mink
Smithsonian Astrophysical Observatory**

American Museum of Natural History, April 13, 2015

Outline

Planet Surfaces: 1973-4 Mars Observations

Stars and Planets: 1976-1990 Occultations

Milky Way: 1985-1990 IRAS and Spacelab 2 IRT

The Universe: 1990-2015 WCS, Plates, and Spectroscopy


American Museum of Natural History, April 13, 2015

Projections

Orthographic: Face-on planet in sky


Mercator: Map of entire planet surface

It All Started With Mars




MIT Vidicon Spectrometer with camera monitoring slit in mirror

Spectrometer slit reprojected across Mercator projection of Mars surface
(Mink, MIT S.M. Thesis 1974)



It All Started With Mars



**Aperture photos projected on observed planet disk
and reprojected onto Mercator projection of Mars**

(Mccord, Huguenin, Mink, and Pieters, Icarus 31, 1977)


Then Came Uranus

Circular No. 3047
 Central Bureau for Astronomical Telegrams
 INTERNATIONAL ASTRONOMICAL UNION
 Postal Address: Central Bureau for Astronomical Telegrams
 Smithsonian Astrophysical Observatory, Cambridge, MA 02138, U.S.A.
 Cable Address: SATELLITES, NEWYORK Telex: 921428
 Telephone: (617) 864-5758

OCCULTATIONS BY URANUS AND (6) HEBE

R. Barrow, Gerard P. Kuiper Airborne Observatory, has relayed word from Perth of successful observations by J. L. Elliot in the southern Indian Ocean of last night's occultation of SAO 158687 by Uranus. A secondary occultation was also observed, this presumably being caused by a small body (not Miranda) in orbit about Uranus. J. Hers reports that heavy rain prevented observations in the vicinity of Johannesburg.

Preliminary reports reaching D. Dunham, Computer Sciences Corporation, suggest that the central line of the occultation of gamma Cet by (6) Hebe passed between 50 and 90 km north of Mexico City. Near the latter point the event lasted 55, beginning on Mar. 5d02h34m54s UT. A 2.5s-duration occultation was observed in Mexico City itself.



Circular No. 3048
 Central Bureau for Astronomical Telegrams
 INTERNATIONAL ASTRONOMICAL UNION
 Postal Address: Central Bureau for Astronomical Telegrams
 Smithsonian Astrophysical Observatory, Cambridge, MA 02138, U.S.A.
 Cable Address: SATELLITES, NEWYORK Telex: 921428
 Telephone: (617) 864-5758

OCCULTATION OF SAO 158687 BY URANUS AND SATELLITE BELT

Amplifying the brief announcement on [IAUC 3047](#), J. L. Elliot reports that several secondary occultations of SAO 158687 on Mar. 10 were observed by E. Dunham, D. Mink and himself from the Kuiper Airborne Observatory and also by R. L. Millis, P. Birch and D. Trout at the Perth Observatory. Both groups independently concluded that these occultations were caused by bodies that are apparently part of a satellite belt about 40 000 km distant from the center of Uranus. The diameters of the satellites range from 100 km to much smaller values. The occultation by Uranus itself was successfully observed from the Airborne Observatory (located at Long. = -90°, Lat. = -50°) and lasted ~ 25 min centered on 21h06m UT. The Uranus occultation did not occur at the Perth Observatory. The secondary occultations took place during an 8-9 min interval around 20h16m UT and during a similar interval around 21h50m UT (although dawn prevented observations of the latter events in Perth).

M. K. V. Bappu, Indian Institute of Astrophysics, cables: "Using the 102-cm reflector at Kavalur (Long. = -5h15m19s.6, Lat. = +12°34'32"), Bhattacharyya and Kuppuswamy found the diminution of SAO 158687 during the occultation by the atmosphere of Uranus to be 0.046 magnitude at an effective wavelength of 7500 Å. Visually and photoelectrically they observed the complete disappearance of the star for 8s.9 beginning at 20h19m15s UT and ascribe this to obscuration by a hitherto unknown satellite of the planet."

Computations by the undersigned show that the asymmetry in the times of the satellite occultations about the main occultation (as observed at the Airborne Observatory) is consistent with the existence of a circular belt in the plane of Uranus' equator. Allowance for foreshortening yields the radii of the inner and outer edges of the belt as 44 000 and 51 000 km, respectively. The Kavalur observation suggests occultation by a 100-km-sized body near the outer edge of the belt. Other observers are urged to examine their records for further evidence of this belt. At Sutherland, times of mid-occultation by the belt would have been 20h22m and 21h59m UT; at Mauritius, 20h23m and 21h54m; at Lembang, 20h20m and 21h46m; at Kyoto, 20h22m; at Helwan, 21h56m. Those wishing to attempt to detect the satellite belt directly are advised that at the present opposition it should be located from 3".5 to 4".0 to the north and south of the center of Uranus and from 2".7 to 3".1 to the east and west; the brightest bodies in it are expected to have $m_v \sim 19$.


1977 March 14

(3048)


Brian G. Marsden

Occultation of SAO 158687 by Uranus and Its Rings
(Eliot, Dunham, and Mink, Nature 261, 328, May 26 1977)

Mapping observability




Geocentric prediction of Uranus 16



Sun down, Uranus up for Uranus 16


Finding Stars to be Occulted



Map showing appulse of the Pluto/Charon system to a star
Note rotation of Charon around Pluto and variations in star position

Predicting Occultations


Centered on kmu 102 $19^{\text{h}}07^{\text{m}}05.365^{\text{s}}$ $-22^{\circ}58'47.1''$
 Uranus from 1 July 1992 0:00 UT every 1 day
 SAO Catalog (SAOra)
 Klemola & Mink Uranus Stars (kmu)



Palomar Sky Survey overlay
 for stars occulted by Uranus

July 8 1992 Occultation of kmu 102 by Uranus epsilon ring
 Observed from IRTF Mauna Kea 3.8m Long= 155 28 15.0 Lat= 19 49 34.0
 13.01-magnitude star at RA= 19h 09m 40.3332s Dec= -22d 54' 34.113"
 (1950) RA= 19h 07m 05.3650s Dec= -22d 58' 47.100"

Closest at 10: 7:25 U.T.
 Radial= 8025.7 km = 0.60"
 Planet= 26145.0 km = 1.95"
 Ring = 51149.3 km = 3.81"
 DE= 60.4757 Pole= 99.7807 deg
 V=0.0017 "/s = 23.05 km/s
 D= 18.525434 AU
 ET - UT = 59.3660 sec DE-130
 Immersion: 9:35:43 U.T.
 P.A.= 273.4 deg. 7/ 8
 R = 44574.2 km = 3.32"
 Phase = 0.0 deg = 100.0%
 Alt= 47 d, Az= 164 d
 Sun alt=-47 d, Az=-17 d
 Sun:179.4 d, Moon: 73.4 d
 V =0.0017 "/s = 23.05 km/s
 VP=0.0019 "/s = 25.88 km/s
 D = 18.525427 A.U.
 Emersion: 10:39:54 U.T.
 P.A.= 73.0 deg. 7/ 8
 R = 45649.7 km = 3.40"
 Phase = 0.0 deg = 100.0%
 Alt= 49 d, Az= 185 d
 Sun alt=-49 d, Az= 4 d
 Sun:179.3 d, Moon: 72.8 d
 V =0.0017 "/s = 23.05 km/s
 VP=-.0019 "/s = -25.89 km/s
 D = 18.525442 A.U.
 Doug Mink 13:33 Mar 18 1992



Sky plane map of Uranus
 ring occultation of KMU102

Predicting Occultations

Venus Occultation of SAO 160149 on January 21, 2003

[Click here for predictions for various cities](#)

Catalog positions at 2003-01-21 11:00 UT

The arcsec column gives the distance from the SAO position

SAO number	RA2000	Dec2000	Mag	Type
160149	16:53:54.416	-19:18:50.17	8.20	G0

PPM number	RA2000	Dec2000	Mag	Type	Arcsec
232288	16:53:54.583	-19:18:48.54	8.60	G0	2.87

Tycho2_num	RA2000	Dec2000	MagB	MagV	Arcsec
6226.02681	16:53:54.550	-19:18:51.04	9.14	8.57	2.08

The following positions are at the catalog epochs


GSC number	RA2000	Dec2000	Mag	Class	Band	N	Arcsec
6226.0268	16:53:54.509	-19:18:50.44	8.51	0	B	1	1.28

GSC-ACTnum	RA2000	Dec2000	Mag	Class	Band	N	Arcsec
6226.0268	16:53:54.552	-19:18:50.15	8.51	0	B	1	1.84

2MASS num.	RA2000	Dec2000	MagJ	MagH	MagK	Arcsec
25.2772036	16:53:54.544	-19:18:50.92	7.399	7.151	7.067	1.93

USNO A2 number	RA2000	Dec2000	MagB	MagR	Plate	Arcsec
0675.16930485	16:53:54.562	-19:18:48.34	9.2	9.9	808	2.62

Finding Charts




[Click for larger map of 5° field around star](#) [Click for larger map of 1° field around star](#)

Predictions for Different Catalog Positions



[Click for larger map of Venus occultation of SAO position](#)



[Click for larger map of Venus occultation of PPM position](#)



[Click for larger map of Venus occultation of Tycho-2 position](#)

IRAS Projections

Infrared Astronomical Satellite (IRAS) Explanatory Supplement (1988)

Edited by C.A. Beichman, G. Neugebauer, H.J. Habing, P.E. Clegg, T.J. Chester

X. The Formats of the IRAS Catalogs and Atlases, D. Extended Emission


Polar: Maps of sky around North and South Poles

Aitoff: Map of entire sky

Sinusoidal: Maps of galactic plane


Gnomonic: Maps of regions of sky

All-Sky IR Mapping from Space



Spacelab 2 Infrared Telescope
(Space Shuttle Challenger, July 1985)

All Sky IR Mapping from Space




IRAS 120 μm
Aitoff All-Sky Projection
1984 Data Release HCON 1

All Sky IR Mapping from Space

```
COMMENT PROJECTION FORMULAE:
COMMENT FORWARD FORMULA; XLON0 IS THE CENTER LONGITUDE OF THE
COMMENT MAP. ARC-SINE AND ARC-COSINE FUNCTIONS ARE REQUIRED.
COMMENT R2D = 45. / ATAN(1.)
COMMENT PIX = 2.
COMMENT RHO = ACOS( COS(XLAT) * COS((XLON-XLON0)/2.) )
COMMENT THETA = ASIN( COS(XLAT) * SIN((XLON-XLON0)/2.) / SIN(RHO) )
COMMENT F = 2. * PIX * R2D * SIN(RHO/2.)
COMMENT SAMPLE = -2. * F * SIN(THETA)
COMMENT XLINE = -F * COS(THETA)
COMMENT IF(XLAT .LT. 0.) XLINE = -XLINE
COMMENT
COMMENT REVERSE FORMULA; XLON0 IS THE CENTER LONGITUDE OF THE MAP.
COMMENT ARC-SINE AND ARC-COSINE FUNCTIONS NEEDED.
COMMENT R2D = 45. / ATAN(1.)
COMMENT PIX = 2.
COMMENT Y = -XLINE / (PIX * 2. * R2D)
COMMENT X = -SAMPLE / (PIX * 2. * R2D)
COMMENT A = SQRT(4.-X*X-4.*Y*Y)
COMMENT XLAT = R2D * ASIN(A*Y)
COMMENT XLON = XLON0 + 2. * R2D * ASIN(A*X/(2.*COS(XLAT)))
COMMENT
COMMENT REFERENCES:
COMMENT IRAS SDAS SOFTWARE INTERFACE SPECIFICATION(SIS) #623-94/NO. SF05
COMMENT ASTRON. ASTROPHYS. SUPPL. SER. 44,(1981) 363-370 (RE:FITS)
COMMENT RECONCILIATION OF FITS PARMS W/ SIS SF05 PARMS:
COMMENT NAXIS1 = (ES - SS + 1); NAXIS2 = (EL - SL + 1);
COMMENT CRPIX1 = (1 - SS); CRPIX2 = (1 - SL)
```

IRAS Aitoff All-Sky Projection Fortran Code in FITS Header

All Sky IR Mapping from IRT



1985 Day 213, Orbit 4, 50,964 0.1-sec frames

Spacelab 2 Infrared Telescope

(Space Shuttle Challenger, July 1985)

Galactic Center from Spacelab 2





FIG. 5.—Contour map of the $2.4 \mu\text{m}$ emission from the Galactic plane region. The contours are spaced logarithmically in 10 steps between 0.67×10^{-10} and $16 \times 10^{-10} \text{ W cm}^{-2} \mu\text{m}^{-1} \text{ sr}^{-1}$.

Linear Projection in Galactic Coordinates


(Kent, Mink, Fazio, Koch, Melnick, Tardiff, Maxson, *ApJS* 78:403-408, 1992)

Galactic Center from Spacelab 2




Linear Projection in Galactic Coordinates
(Mink, August 1990, unpublished)

All-Sky Maps meet Catalogs




All-Sky Maps meet Catalogs




Space Telescope Guide Star Catalog, Galactic Plane, Aitoff Projection
(Mink, D.J. 1994. In *Astronomical Data Analysis Software and Systems III*, A.S.P. Conference Series, Vol. 61, 1994, Dennis R. Crabtree, R.J. Hanisch, and Jeannette Barnes, eds., p. 191)

Galileo's Telescope Expands the Sky

Rome, Italy
April 14, 1611 7:00 PM
Sky to 6th magnitude
(Naked eye)




Rome, Italy
April 14, 1611 7:00 PM
Sky to 8th magnitude
(Galileo's telescope)



Polar projections centered on local apex in Rome

Owen Gingerich will present a keynote address at a conference sponsored by the American Academy of Rome celebrating the moment in 1611 when Galileo Galliei proudly presented the "telescope" to the intelligentsia of Rome... the Academy asked Paine Professor of Astronomy and director of the Harvard-Smithsonian Center for Astrophysics Irwin Shapiro and his Center colleague Douglas Mink to produce a map of the stars as they appeared over Rome on the night of April 14, 1611. For their efforts, Shapiro and Mink received a Jeroboam of champagne; Gingerich, however, got a trip to Rome. (Harvard Gazette, April 10, 1997)

Onto the World Wide Web




The screenshot shows a Netscape browser window with the following content:

File Options Navigate Annotate Help

Document Title: SAO/TDC Home Page

Document URL: file:///localhost/cfa165/www/TDC.html

 **Telescope**
SMITHSONIAN ASTROPHYSICAL OBSERVATORY
Data Center

The Smithsonian Astrophysical Observatory (SAO) is part of the Harvard-Smithsonian Center for Astrophysics in Cambridge, Massachusetts. The SAO/TDC creates and maintains software to process and archive data from optical and infrared telescopes in Harvard, Massachusetts and on Mt. Hopkins in Arizona.

The SAO TDC distributes several pieces of software:

- [RVSAO](#): an IRAF package for finding radial velocities from spectra
- [RGSC](#): a program for searching the Hubble Space Telescope Guide Star Catalog.
- [SKYMAP](#): a program for mapping star catalogs onto the sky

The staff of the TDC are:

- [Doug Mink](#), software developer
- [Mike Kurtz](#), software philosopher

Astronomical Software Elsewhere

Other Astronomical Resources

Navigating the Internet

A useful introduction to the **World-Wide Web (WWW)** is available from NASA/Goddard.

Last updated 12 November 1993

[Back](#) [Forward](#) [Home](#) [Reload](#) [Open...](#) [Save As...](#) [Clone](#) [New Window](#) [Close Window](#)

Documentation on the World Wide Web



The screenshot shows a web browser window with the address bar containing `tdc-www.harvard.edu/software/skymap/`. The browser's address bar includes navigation icons (back, forward, refresh) and a search icon. Below the address bar, there are tabs for 'Apps', 'Five Labs | Jessica', and 'ADSLABS'. The main content area has a light blue background with a starry pattern. On the left, there is a logo for SKYMAP, which consists of two blue stars flanking the word 'SKYMAP' in a blue oval. On the right, there is a logo for the Telescope Data Center, featuring a yellow sun icon and the text 'Telescope Data Center' and 'SMITHSONIAN ASTROPHYSICAL OBSERVATORY'. The main text describes SKYMAP as an astronomical mapping program written in Fortran and C for Unix workstations by Doug Mink of the Smithsonian Astrophysical Observatory Telescope Data Center. It lists various options and catalogs, including GSC, USNO, SAO, WCSTools, and programs like rgsc and star. The text is followed by a list of links for 'Manual', 'Examples', 'Commands', 'Installation', 'Parameters', 'Catalogs', 'Reference', 'Notes', and 'Versions'. At the bottom, it states 'Last updated 3 April 2003 by Doug Mink, dmink@cfa.harvard.edu'.

tdc-www.harvard.edu/software/skymap/

Apps Five Labs | Jessica ADSLABS

 SKYMAP is an astronomical mapping program written in Fortran and C for Unix workstations by [Doug Mink](#) of the [Smithsonian Astrophysical Observatory Telescope Data Center](#). If you just need positions, several other options are available. The [WCSTools](#) package contains [C programs](#) which can search the [GSC](#), [USNO](#), and [SAO](#) catalogs, among [others](#). The obsolete programs, [rgsc](#) for the Guide Star Catalog and [star](#) and its variants for other catalogs, use the same Fortran and C code as **skymap**.

 **Telescope Data Center**
SMITHSONIAN ASTROPHYSICAL OBSERVATORY

[Manual](#)

[Examples](#) [[Grid](#)] [[Guide Stars](#)] [[New Field](#)] [[Field from catalog](#)]

[Commands](#) [[Command Line](#)] [[Menu](#)] [[Cursor](#)]

[Installation](#)

Parameters: [[Dictionary](#)] [[Format](#)]

[Catalogs](#) [[ASCII](#)] [[Binary](#)]

Reference [D.Mink\(1993\)](#), [ADASS II](#) [[full text](#)]

[Notes](#)

[Versions](#)

Last updated 3 April 2003 by [Doug Mink](#), dmink@cfa.harvard.edu

Digitized Sky Survey Projection

AURA (1993-1994), "The Digitized Sky Survey" CDROM Manual


Doggett, J. (1997), <http://gsss.stsci.edu/Software/GetImage/GetImage.htm>

Mink, Jessica (1999), "WCSTools: Image Astrometry Toolkit", <http://tdc-www.harvard.edu/software/wcstools/>


PLTLABEL= 'E1356	' /Observatory plate label	PPO1 = 0.000000000000E+00	/Orientation Coefficients
PLATEID = '08MC	' /GSSS Plate ID	PPO2 = 0.000000000000E+00	/
REGION = 'XE429	' /GSSS Region Name	PPO3 = 1.7747471555000E+05	/
DATE-OBS= '23/03/55	' /UT date of Observation	PPO4 = 0.000000000000E+00	/
UT = '06:02:00.00	' /UT time of observation	PPO5 = 0.000000000000E+00	/
EPOCH = 1.9552226562500E+03	/Epoch of plate	PPO6 = 1.7747471555000E+05	/
PLTRAH = 10	/Plate center RA	AMDX1 = 6.7241844402360E+01	/Plate solution x coefficients
PLTRAM = 7	/	AMDX2 = 3.9757845495110E-01	/
PLTRAS = 5.5528480000000E+01	/	AMDX3 = -2.0498717200880E+02	/
PLTDECSN= '+'	' /Plate center Dec	AMDX4 = -1.3607216767070E-05	/
PLTDECD = 17	/	AMDX5 = -2.2201873529570E-05	/
PLTDECM = 17	/	AMDX6 = 7.4284599162830E-07	/
PLTDECS = 3.8380860000000E+01	/	AMDX7 = 0.000000000000E+00	/
EQUINOX = 2.000000000000E+03	/Julian Reference frame equinox	AMDX8 = 1.9162087720540E-06	/
EXPOSURE= 5.000000000000E+01	/Exposure time minutes	AMDX9 = -9.2146076767620E-10	/
BANDPASS= 8	/GSSS Bandpass code	AMDX10 = 2.1089546241680E-06	/
PLTGRADE= 1	/Plate grade	AMDX11 = -9.3945135632070E-08	/
PLTSCALE= 6.720000000000E+01	/Plate Scale arcsec per mm	AMDY1 = 6.7256622034650E+01	/Plate solution y coefficients
SITELAT = '+33:24:24.00	' /Latitude of Observatory	AMDY2 = -3.9844579471320E-01	/
SITELONG= '-116:51:48.00	' /Longitude of Observatory	AMDY3 = -6.8591056129270E+01	/
TELESCOP= 'Palomar 48-inch Schmidt'	/Telescope where plate taken	AMDY4 = -1.3176449798960E-05	/
CNPIX1 = 10748	/X corner (pixels)	AMDY5 = -7.8391468151820E-06	/
CNPIX2 = 2023	/Y corner	AMDY6 = -7.4802178840710E-07	/
DATATYPE= 'INTEGER*2	' /Type of Data	AMDY7 = 0.000000000000E+00	/
XPIXELSZ= 2.528445000000E+01	/X pixel size microns	AMDY8 = 1.8834016180180E-06	/
YPIXELSZ= 2.528445000000E+01	/Y pixel size microns	AMDY9 = -1.9452422448560E-07	/
		AMDY10 = 2.1574073462190E-06	/
		AMDY11 = -1.6009508926300E-08	/

Digitized Sky Survey Projection

*Mink, Jessica (1999), "WCSTools: Image Astrometry Toolkit"
, <http://tdc-www.harvard.edu/software/wcstools/>*



**DS9 display of DSS image
with previous header**



**DS9 display of same DSS image with regions
generated by WCSTools imcat program
which remotely accessed the GSC2 catalog**

AIPS Projections

The eight most commonly-used projections of classic AIPS may be computed using the *worldpos* and *worldpix* subroutines written by Bill Cotton and Eric Greisen of NRAO:

SIN: Orthographic projection

TAN: Tangent plane projection

ARC: Zenithal equidistant projection

NCP: North celestial pole projection

GLS: Sanson-Flemsteed sinusoidal projection

MER: Mercator projection

AIT: Hammer-Aitoff equal area all-sky projection


STG: Stereographic projection
(zenithal orthomorphic)

FITS-WCS Projections

*Calabretta, M.R., & Greisen, E.W., (2002), "Representations of celestial coordinates in FITS",
Astronomy & Astrophysics, 395, 1077-1122.*

AZP: Zenithal (Azimuthal) Perspective	COD: COnic equiDistant
SZP: Slant Zenithal Perspective	COE: COnic Equal area
TAN: Gnomonic = Tangent Plane	COO: COnic Orthomorphic
SIN: Orthographic/synthesis	BON: Bonne
STG: Stereographic	PCO: Polyconic
ARC: Zenithal/azimuthal equidistant	SFL: Sanson-Flamsteed
ZPN: Zenithal/azimuthal PolyNomial	PAR: Parabolic
ZEA: Zenithal/azimuthal Equal Area	AIT: Hammer-Aitoff equal area all-sky
AIR: Airy	MOL: Mollweide
CYP: CYlindrical Perspective	CSC: COBE quadrilateralized Spherical Cube
CAR: Cartesian	QSC: Quadrilateralized Spherical Cube
MER: Mercator	TSC: Tangential Spherical Cube
CEA: Cylindrical Equal Area	NCP: North celestial pole (special case of SIN)
COP: COnic Perspective	GLS: GLobal Sinusoidal (Similar to SFL)

Fitting a WCS using WCSTools




USNO-B1.0 Catalog plotted over image using telescope pointing

```
NAXIS = 2 / NUMBER OF AXES
NAXIS1 = 1024
NAXIS2 = 1024
RA = '16:15:56' /MEAN RA
DEC = '31:25:10' /MEAN DEC
EPOCH = 2000 /MEAN EPOCH
DATE-OBS= '1995-03-06T10:04:50' /UT DD/MM/YY AT END OF EXPOSURE
EXPTIME = 240.00 /INTEGRATION TIME, SECONDS
HJDN = 2449871.91686 /HELIOCENTRIC JULIAN DAY AT MIDDLE OF EXPOSURE
SECPIX = 0.652 /SEC OF ARC SPATIAL PIXEL , SET BY PARAMETERS
```

FITS header with limited WCS keywords from telescope

Fitting a WCS using WCSTools



USNO-B1.0 Catalog plotted over image after imwcs WCS fit

```
NAXIS = 2 / NUMBER OF AXES
NAXIS1 = 1024
NAXIS2 = 1024
RA = '16:15:56.591'
DEC = '+31:25:07.35'
EPOCH = 2000
RADECSYS= 'FK5'
EQUINOX = 2000
CRPIX1 = 516.9970
CRPIX2 = 513.4151
CD1_1 = -0.000178936537
CD1_2 = 0.000000555651
CD2_1 = 0.000000278868
CD2_2 = 0.000178740421
WCSRFCAT= 'ub1'
WCSIMCAT= 'testx90n.sex'
WCSMATCH= 147
WCSNREF = 217
WCSTOL = 2.5000
CTYPE1 = 'RA---TAN'
CTYPE2 = 'DEC--TAN'
CRVAL1 = 243.985795481
CRVAL2 = 31.418709691
SECPIX1 = 0.6442
SECPIX2 = 0.6435
WCSSEP = 0.257
```

FITS header with WCS keywords after WCS fit using imwcs

Finding Stars for WCSTools

WCSTools supports several ways to find the star-like objects in an image


- **WCSTools IMSTAR task (also built into IMWCS)**
- **IRAF DAOFIND task**
(its X,Y,Magnitude is WCSTools standard position format)
- **SExtractor**
(output formattable to WCSTools standard)

More Catalogs

These catalogs are available and supported by [SAO/TDC search and mapping software](#).


Catalog or Format	No. Stars	Bytes	Region Search	Image Search	Mapping
USNO-B1.0 Catalog	1,036,366,767	83,614,060,960	sub1 (scat)	imub1 (imcat)	
GSC II Catalog (2.2.01)	998,402,801	>80,000,000,000	sgsc2 (scat)	imgsc2 (imcat)	
2MASS Point Source Catalog	470,992,970	31,775,592,000	stm2 (scat)	imtm2 (imcat)	
2MASS Extended Source Catalog	1,647,599	11,533,193	stmx (scat)	imtmx (imcat)	
USNO UCAC3 Catalog	100,766,420	8,536,270,559	sucac3 (scat) ,	imucac3 (imcat)	
USNO UCAC2 Catalog	48,366,996	2,128,147,841	sucac2 (scat) ,	imucac2 (imcat)	
GSC-ACT Catalog	25,541,952	1,231,787,520	sgsca (scat)	imgsca (imcat)	
SDSS Photometry Catalog	53 million (DR1)		?? ssdss (scat)	imscdss (imcat)	
HST Guide Star Catalog	25,541,952	1,231,787,520	sgsc (scat)	imgsc (imcat)	skymap
Tycho-2 Catalog	2,539,913	528,721,576	sty2 (scat)	imty2 (imcat)	
USNO/Hipparcos ACT Catalog	988,758	318,380,076	sact (scat)	imact (imcat)	
SKY2000 Catalog	299,167	11,368,374	ssky2k (scat)	imsky2k (imcat)	skymap
PPM Catalog	378,910	22,734,656	sppm (scat)	imppm (imcat)	skymap
SAO Catalog	258,996	15,539,876	ssao (scat)	imsao (imcat)	skymap
IRAS Point Source Catalog	245,889	7,376,698	siras (scat)	imiras (imcat)	skymap
Hipparcos Catalog	118218	4492312	ship (scat)	imhip (imcat)	skymap
Yale Bright Star Catalog	3256	291548	sbsc (scat)	imbsc (imcat)	skymap
Starbase tab-delimited ASCII	varies	varies	scat	imcat	
TDC Space-Delimited ASCII	varies	varies	scat	imcat	skymap
TDC Binary	varies	varies	scat	imcat	skymap
USNO-A2.0 Catalog	526,280,881	6,315,370,572	sua2 (scat)	imua2 (imcat)	skymap
USNO-SA2.0 Catalog	55,368,239	664,418,868	susa2 (scat) ,	imusa2 (imcat)	
USNO-A1.0 Catalog	488,006,860	5,856,082,320	sua1 (scat)	imua1 (imcat)	skymap
USNO-SA1.0 Catalog	54,787,624	657,451,488	susa1 (scat)	imusa1 (imcat)	skymap
USNO J-1.0 Catalog	19,911,514	238,938,168	sujc (scat)	imujc (imcat)	

Testing Catalog Accuracy



216 1x1 degree fields from the 8K array on the KPNO 36-inch telescope cover half of the CfA Century survey, 50 degrees across the sky





Testing Catalog Accuracy



The 2MASS Point Source Catalog plotted over one of the 1728 test images

Testing Catalog Accuracy




D. J. Mink, W. R. Brown, M. J. Kurtz (2004) A Comparison of Large All-Sky Catalogs
Astronomical Data Analysis Software and Systems XIII, F. Ochsenbein, M. Allen,
and D. Egret, eds. ASP Conference Series, Vol. 314, p. 141



Differences between catalogs and WCS fits to 1726 images

Testing Catalog Accuracy

D. J. Mink (2010) Unpublished



Differences between catalogs and WCS fits to 1726 images


Accommodating image distortions in WCSTools

WCSTools supports several methods of fitting distortions to images as this seems not to be standardizable

- Digitized Sky Survey Plate Model
- IRAF TNX and ZPX projections with polynomial distortion
- Spitzer/STScI polynomial distortion model
- SWARP polynomial distortion model

Putting Positions on Harvard's Plates


Mink, D.; Doane, A.; Simcoe, R.; Los, E.; Grindlay, J. (2006). "The Harvard Plate Scanning Project" in Virtual Observatory: Plate Content Digitization, Archive Mining and Image Sequence Processing, iAstro workshop, Sofia, Bulgaria, 2005 proceedings, Sophia: Heron Press Ltd.



From hand-written cards and logbooks

Putting Positions on Harvard's Plates

Mink, D.; Doane, A.; Simcoe, R.; Los, E.; Grindlay, J. (2006). "The Harvard Plate Scanning Project" in Virtual Observatory: Plate Content Digitization, Archive Mining and Image Sequence Processing, iAstro workshop, Sofia, Bulgaria, 2005 proceedings, Sophia: Heron Press Ltd.




From three floors of cabinets of glass plates

Putting Positions on Harvard's Plates



File Edit View History Bookmarks Tools Help
edu/cgi-bin/plates/mcplate?rfn=21475&dx= per pipe
python h... Full Text ... Pipes (Pr... Perf File L... MC Serie... MC ... x mcpart (...)

 **Telescope Data Center**
SMITHSONIAN ASTRONOMICAL OBSERVATORY

[Click here for 1/32nd scale FITS image with WCS](#)
[Click here for 1/16th scale FITS image with WCS](#)
[Click on image for 1000 x 1000 JPEG image.](#)

Harvard Plate
Stacks
MC Series Plate
MC21475

[A Series \(about\)](#)
[MC Series \(about\)](#)
[MF Series \(about\)](#)


Plate	RA2000	Dec2000	Exp	Epoch	Note
21475	08:39:44.900	+19:44:52.60	25.00	1925-05-22T00:00	RY_Cancri

From glass to bytes on home-built scanner

Putting Positions on Harvard's Plates



M44 in Plate MC21438



**M44 in Plate MC21438
with Tycho 2 Catalog stars marked**

Zoom in and overplot stars using WCS

Putting Positions on Harvard's Plates



100,000th Plate Scanned, April 7, 2015

sethead sets values of keywords in FITS Headers

Each keyword should be followed by an equal sign and the value to which it is to be set

Values which are all numeric are assumed to be numbers and are aligned as such within the header.

A list of filenames may be used by prefacing the name of the file containing the list with a @.

Multiple FITS extension headers may be changed at once using -x [range of extension numbers]

Change the right ascension and declination of a FITS image to a different epoch.

Before:

```
RA    = '9:51:19.45'    /MEAN RA
DEC   = '69:15:26.42'  /MEAN DEC
EPOCH = 1950           /MEAN EPOCH
```

```
$ sethead -nvkr X ra='09:55:25.177' dec='+69:01:13.72' epoch=J2000 test.fits
SETHEAD WCSTools 3.9.1, 24 March 2015, Jessica Mink (jmink@cfa.harvard.edu)
Set Header Parameter Values in FITS image file test.fits
RA = 09:55:25.177
DEC = +69:01:13.72
EPOCH = 2000
teste.fits: rewritten successfully.
```

After:

```
XRA   = '9:51:19.45'    /MEAN RA
XDEC  = '69:15:26.42'  /MEAN DEC
XEPOCH = 1950          /MEAN EPOCH
```

...

```
RA    = '09:55:25.177'
DEC   = '+69:01:13.72'
EPOCH =                2000
SETHEAD = 'SETHEAD 2.5 1998-09-01 13:31 RA, DEC, EPOCH updated'
```

gethead extracts information from FITS headers

Keyword names may be entered in either upper or lower case

Tab-separated table output, with column headers, is an option

Afile containing a list of filenames may be used by prefacing it with a @.

Multiple parameters from list of FITS files

Get the image sizes from the NAXIS, NAXIS1, and NAXIS2 header keywords from a list of FITS and IRAF files, printing the output in tab table format:

```
$ gethead -th @fits.list naxis naxis1 naxis2
FILENAME          NAXIS    NAXIS1   NAXIS2
-----
0083.19083010-0706459.fits      2          2720    161
hiptest.fits          2          600     600
test.fits             2          2720    161
test_fabien.fits      2          2080    2048
testbin.fits          2          765     510
testbinf.fits         2          680     450
testbinfg10x10.fits    2          765     510
testbinfg20x20.fits    2          765     510
testbinfg40x40.fits    2          680     450
webccd-1.fits         2          680     450
webccd-2.fits         2          765     510
```

